

SOFIA follow-ups of ATLASGAL massive clumps


Friedrich Wyrowski, Rolf Güsten, Karl Menten,
Helmut Wiesemeyer & Silvia Leurini

MPIfR Bonn





Massive clump evolution

Infall is a fundamental process in SF!

G023.2056-0.3772 IRAC + 870 μ m Contours



?



Outline



- The ATLASGAL survey
- Molecular line follow-ups
- How to probe infall
- Previous ammonia results
- New ammonia observations
- The extended sample
- Comparing ammonia with other high density probes
- New modeling of profiles
- APEX-SOFIA synergies

Overview of the ATLASGAL Survey

Submillimetre Emission: Structure of the Dust


- APEX 870 μm continuum survey of the inner Galactic plane ($300^\circ < l < 60^\circ$, $|b| < 1.5^\circ$)
- APEX: 12-m Single-dish Submillimetre Telescope located on the Chajnantor Plateau
- LABOCA: Large APEX BOlometer CAmera (MPIfR)
- 295 elements at 870 μm
- FoV = $11'$, an angular resolution of $\sim 19''$ and sensitivity of ~ 60 mJy/beam

Complementary to Herschel/HiGAL
&
Pathfinder to ALMA MSF science




Overview of the ATLASGAL Survey

Submillimetre Emission: Structure of the Dust



Molecular line follow ups

- Dust continuum is important but molecular line information is indispensable !
- Effelsberg/Parkes NH_3 (Wienen+'12, '15):
 - Kinematic distances, temperatures
- IRAM 30m/ATNF-Mopra/APEX (Wyrowski/Csengeri):
 - 3 & 0.85mm line surveys: Physical & chemical conditions (CO , Giannetti+2014)
- Herschel/HIFI: 100 ATLASGAL sources currently observed in water lines \rightarrow ATLASGAL water legacy (Wyrowski+; IRAS17233: Leurini+2014)
- MALT90 (Jackson+2013): Mopra Galactic Plane Survey of high density regions. large program:
 - \rightarrow about 2000 ATLASGAL clumps mapped at 3mm
- SEDIGSM (Schuller+): APEX $^{13}\text{CO}(2-1)$ survey of southern Galactic Plane




König+2015

ATLASGAL @ SOFIA

concerted effort on several fronts












- Massive clumps selected in several GMCs (e.g. Wienen+2015)
- Observe multiple sources without changing flight directions
- Multi-semester project
- → statistical significant sample covering range of evolutionary stages
- Cooling budget: OI/CII (e.g. Leurini talk)
- High- J CO: (11-10)/(16-15)
- Ammonia infall study



Wienen+2015: HI+870 μ +¹³CO

SOFIA follow-ups (CO teaser)

G014.1944-0.1939 WISE + 870 μ m Contours





Search for infall

I: Blue-skewed profiles

Needs excitation gradient, right tau

II: red-shifted absorption

Needs high critical density, central continuum



The story so far:

Ammonia@1.8THz

Wyrowski+2012

- 3 absorption line detections in science verification
- All redshifted with respect to v_{sys}
- $\tau \sim 1$

Table 2. Line parameters from Gaussian fits to the NH_3 lines. Nominal fit errors are given in brackets. In addition, the velocity of C^{17}O (3-2) lines observed with the APEX telescope are given.

Source	T_{peak} (K)	Δv (km s^{-1})	$v_{\text{LSR}}^{\text{NH}_3}$ (km s^{-1})	$v_{\text{LSR}}^{\text{C}^{17}\text{O}}$ (km s^{-1})
W43-MM1	-0.96 (0.22)	5.3 (0.8)	99.7 (0.4)	97.65 (0.06)
G31.41+0.31	-1.18 (0.29)	3.7 (0.8)	99.4 (0.4)	97.02 (0.04)
G34.26+0.15	-3.38 (0.56)	5.5 (0.6)	61.2 (0.3)	58.12 (0.03)




Fig. 2. NH_3 spectra of the observed sources. Results of Gaussian fits to the line are overlaid in green. The systemic velocities of the sources, determined using C^{17}O (3-2) are shown with dotted lines.

G34: comparison to VLA absorption





Fig. 3. G34.26+0.15 SOFIA NH₃ spectrum compared with the VLA NH₃ (2,2) spectrum taken from Gómez et al. (2000) which was integrated over the region which shows absorption. A two-component hyperfine fit to the (2,2) spectrum is shown in green.

Infall Results

- **3 clear detections of Ammonia line-of-sight infall** consistent with results from cm-absorption and/or blue-skewed emission profiles
- More direct probe of infall that **can be extended to earlier stages** of SF without cm background continuum and cases where other species are depleted
- **Infall rates of $3-10 \times 10^{-3} M_{\odot}/\text{yr}$** (if spherical)
- **Next step: extend to more sources and stages, in particular earlier ones**

SOFIA Observations

- GT and cycle 1 science flights
- **GREAT:**
 - L1, various lines
 - NH_3 3_{2+} -2_{2-} 1810.379 GHz LSB
 - AFFTS/XFFTS: 1.5/2.5 GHz
 - Chopped observations of 9 sources



Cycle I: a) continuation to Infrared dark clouds






Figure 2: IRDC G23.21-0.38: (a) ATLASGAL 870 μm dust continuum as contours on GLIMPSE 8 μm MIR emission in color, SOFIA beam in red. (b) APEX HNC (4-3) spectrum of this clump with systemic velocity indicated.

Cycle I: b) filling in further stages:

- **G35.20-0.74**: submm brightest, northern massive young stellar object, fulfilling Lumsden+ MSX color criteria
- **G327.3/G351.58**: hot cores/ultracompact HII regions with high luminosity (up to $2 \times 10^5 L_{\odot}$)

New SOFIA results: sample



König+2015

No/too weak SOFIA 1.8THz continuum ! \rightarrow 572 GHz

New SOFIA results: continuum




Fig. 2. Comparison of GREAT continuum levels with PACS 160 μm flux densities with nominal 20% errors.




Fig. 3. Example of constraining the radial physical structure with the help of the ATLASGAL submm dust continuum radial profiles.

New SOFIA results: lines

Wyrowski et al.: Infall through the evolution of high-mass star forming clumps





Fig. 1. NH_3 $3_{2+} - 2_{2-}$ spectra of the observed sources. Results of Gaussian fits to the line are overlaid in green. The systemic velocities sources, determined using C^{17}O (3–2), are shown with dotted lines.

- 5 new redshifted absorption with shifts of 0.2 – 1.6 km/s with respect to C^{17}O
- 1 source dominated by outflow (G5.89), several blue wings
- 2 sources with blue shifted absorption

How consistent are different probes ?

	nh3	hco+	1-0 hnc	cs	4-3 hnc	hcn	7-6 hco+	cs
G327.29-0.58	o	o	o		o	o	o	+
G351.58-0.35	++	++	++		o/r.a.	o/ra	o/ra	-
G23.21-0.38	++	--	--	--	++	o	++	o
G34.41 irdc43	++	--	--	o	o	o	+	o
G35.19-0.74	+	++	o	--	o	--	++	o
g31.41	++	--	-	++	++	++	++	o
g34.26	++	r.a.	r.a.	++	++	++	++	+
w43mm1/g30.82	++	+	++		++	++	++	o

→ Ammonia and HCO⁺ (4-3) show best correspondence





Modeling

- Fit dust continuum (ATLASGAL) with **density power law** ($n \sim r^\alpha$
 $\alpha = 1.5 - 2.2$)
- **Temperature structure** dictated by inner heating source
(luminosities known from SED fits, König+2015)
- → **Adjust ammonia abundance and velocity structure in spherical RATRAN models**
- Modelling of new sources results in **infall with fractions of free-fall of 5 – 25 %**. The ammonia abundance are in the range of $0.2 - 2 \times 10^{-8}$.
- But how consistent are models of NH_3 and HCO^+ ?

New SOFIA modeling

Outflow component

HCO⁺ usually probing
additional outflow component
→ RATRAN modification of
Mottram+2013




Additional parameter:

- outflow widths/strength
- HCO⁺ abundance


New SOFIA modeling

Outflow component




Also this clump shows consistent results!

But many cases do not work with this simple geometry.
Complicated outflows?
Additional low density outer layer (Lopez-Sepulcre+2010)?




Herschel G34.26 results

Hajigholi+2015



- Different excitation traces different v
- SOFIA opportunities:
 - GS 572 GHz @ 90% transmission
 - 1214 GHz (201-100, 211-110) @ 65%
 - 2355 GHz (4-3 lines) @ 63%



APEX-SOFIA synergies

- Mid vs. High J CO
- Blue-skewed self-absorbed high density probes vs. red-shifted absorption studies
- CO/Cl cooling vs. CII/OI
- Complex molecules vs. hydrides
- Similar beamsizes in APEX submm windows and with SOFIA THz RX
- Imaging: CHAMP+/LASMA vs. UPGREAT



Summary & Outlook

- Infall on clump scales ubiquitous through wide range of evolutionary stages
- Ammonia and HCO^+ (4-3) show best correspondence but HCO^+ stronger affected by outflows
- Continue filling in stages (populating the M-L diagram)
- Study infall across clumps (upGREAT)
- Add additional lines to cover larger excitation range (new single pixel RXs)